

ETC3550/ETC5550 Applied forecasting

Week 7: ARIMA models

af.numbat.space



Outline

1 ARIMA models

2 Stationarity

3 Differencing

ARIMA models

- AR:** autoregressive (lagged observations as inputs)
- I:** integrated (differencing to make series stationary)
- MA:** moving average (lagged errors as inputs)

An ARIMA model is rarely interpretable in terms of visible data structures like trend and seasonality. But it can capture a huge range of time series patterns.

Outline

1 ARIMA models

2 Stationarity

3 Differencing

Stationarity

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

Stationarity

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

Transformations help to **stabilize the variance**.

For ARIMA modelling, we also need to **stabilize the mean**.

Outline

1 ARIMA models

2 Stationarity

3 Differencing

Differencing

Differencing

- Differencing helps to **stabilize the mean**.
- First differencing: *change* between consecutive observations: $y'_t = y_t - y_{t-1}$.
- Seasonal differencing: *change* between years: $y'_t = y_t - y_{t-m}$.

Your turn

- 1 Does differencing make the Closing stock price series stationary for Amazon and Apple stocks?
- 2 What sorts of transformations and differencing are needed to make the Gas series from `aus_accommodation` stationary?

Automatic differencing

Statistical tests to determine the required order of differencing.

- 1 Augmented Dickey Fuller test: null hypothesis is that the data are **non-stationary** and non-seasonal.
- 2 Kwiatkowski-Phillips-Schmidt-Shin (KPSS) test: null hypothesis is that the data are **stationary** and non-seasonal.

Seasonal strength

STL decomposition: $y_t = T_t + S_t + R_t$

Seasonal strength $F_s = \max\left(0, 1 - \frac{\text{Var}(R_t)}{\text{Var}(S_t + R_t)}\right)$

If $F_s > 0.64$, do one seasonal difference.

Automatic differencing

Statistical tests to determine the required order of differencing.

- 1 Augmented Dickey Fuller test: null hypothesis is that the data are **non-stationary** and non-seasonal. H_0 : non-stationary
- 2 Kwiatkowski-Phillips-Schmidt-Shin (KPSS) test: null hypothesis is that the data are **stationary** and non-seasonal. H_0 : stationary

Seasonal strength

STL decomposition: $y_t = T_t + S_t + R_t$

Seasonal strength $F_s = \max\left(0, 1 - \frac{\text{Var}(R_t)}{\text{Var}(S_t + R_t)}\right)$

If $F_s > 0.64$, do one seasonal difference.

Your turn

- 3 Do the unit root tests for the Gas series from `aus_accommodation`. Do they give the same numbers of difference as you chose?