

# **ETC3550/ETC5550 Applied forecasting**

## **Week 7: ARIMA models**

**`af.numbat.space`**



# Outline

1 ARIMA models

2 Stationarity

3 Differencing

# ARIMA models

- AR:** autoregressive (lagged observations as inputs)
- I:** integrated (differencing to make series stationary)
- MA:** moving average (lagged errors as inputs)

An ARIMA model is rarely interpretable in terms of visible data structures like trend and seasonality. But it can capture a huge range of time series patterns.

# Outline

1 ARIMA models

2 Stationarity

3 Differencing

# Stationarity

## Definition

If  $\{y_t\}$  is a stationary time series, then for all  $s$ , the distribution of  $(y_t, \dots, y_{t+s})$  does not depend on  $t$ .

# Stationarity

## Definition

If  $\{y_t\}$  is a stationary time series, then for all  $s$ , the distribution of  $(y_t, \dots, y_{t+s})$  does not depend on  $t$ .

Transformations help to **stabilize the variance**.

For ARIMA modelling, we also need to **stabilize the mean**.

# Outline

1 ARIMA models

2 Stationarity

3 Differencing

# Differencing

## Differencing

- Differencing helps to **stabilize the mean**.
- First differencing: *change* between consecutive observations:  $y'_t = y_t - y_{t-1}$ .
- Seasonal differencing: *change* between years:  $y'_t = y_t - y_{t-m}$ .



# Your turn

- 1 Does differencing make the Closing stock price series stationary for Amazon and Apple stocks?
- 2 What sorts of transformations and differencing are needed to make the Gas series from `aus_accommodation` stationary?

# Automatic differencing

Statistical tests to determine the required order of differencing.

- 1 Augmented Dickey Fuller test: null hypothesis is that the data are **non-stationary** and non-seasonal.
- 2 Kwiatkowski-Phillips-Schmidt-Shin (KPSS) test: null hypothesis is that the data are **stationary** and non-seasonal.

## Seasonal strength

STL decomposition:  $y_t = T_t + S_t + R_t$

Seasonal strength  $F_s = \max\left(0, 1 - \frac{\text{Var}(R_t)}{\text{Var}(S_t + R_t)}\right)$

If  $F_s > 0.64$ , do one seasonal difference.

# Automatic differencing

Statistical tests to determine the required order of differencing.

- 1 Augmented Dickey Fuller test: null hypothesis is that the data are **non-stationary** and non-seasonal.  $H_0$ : non-stationary
- 2 Kwiatkowski-Phillips-Schmidt-Shin (KPSS) test: null hypothesis is that the data are **stationary** and non-seasonal.  $H_0$ : stationary

## Seasonal strength

STL decomposition:  $y_t = T_t + S_t + R_t$

Seasonal strength  $F_s = \max\left(0, 1 - \frac{\text{Var}(R_t)}{\text{Var}(S_t + R_t)}\right)$

If  $F_s > 0.64$ , do one seasonal difference.

# Your turn

- 3 Do the unit root tests for the Gas series from `aus_accommodation`. Do they give the same numbers of difference as you chose?